

Impact of heat stress on poultry: How to reduce the impact on birds and prevent losses of performances during the summer period?

Half of the birds reared worldwide are estimated to suffer from the consequences of heat stress during the summer period. The first visible sign is an increase in the water: feed ratio, however recent research has also shown that excessively hot conditions have an impact on the gut structure and its permeability. Reduced feed consumption and decreased intestinal integrity and function result in the decreased energy intake, animal discomfort and the reduced performance parameters associated with summer months. An effective building temperature control systems is a primary step toward managing this problem but it can be sensitive in case of sudden or excessive changes in temperature. The temperature management can therefore be actively supported by natural nutritional feeding strategies.


“Leaky gut” phenomenon in heat-stressed poultry

Heat is produced by the body metabolism, which includes maintenance, growth and egg production. Birds become 'heat stressed' when they have difficulty achieving a balance between body heat production and body heat loss. This can occur in all ages, and all types of poultry. Birds have three ways of managing their heat loss to stay in the 'thermo-neutral zone'. One of the methods consists of a natural diversion of the blood flow from the viscera to their external surface (wings, skin...) in order to increase the heat exchange rate with the surrounding environment and therefore decrease body temperature. The associated lack of digestive tract blood irrigation progressively leads to gut mucosa depletion, associated with its membrane permeability and its inability to secrete digestive enzymes. This phenomenon already well proven is referred to as “leaky gut”.

Natural ingredients to alleviate and prevent heat stress

Based on the scientific explanations of leaky gut, Pancosma SA, the Swiss additive manufacturer, recently evaluated the efficacy of one of its leading products, known to be a powerful gut protector. The product XTRACT® 6930 is already well known for its multifunctional positive effects on digestive secretions, reduced inflammation and antioxidant properties. It was therefore tested in a heat stress controlled trial at Harper Adams university (UK ; supervised by Dr. Pirgozliev). A total of 36 male Ross 308 broiler chickens were reared from hatching until 20 days old in a common floor pen and under normal recommended temperature. At day 21, the temperature was increased from 21°C to 35°C, simulating a violent environmental heat stress, this continued until the trial finished on day 35. The effects of 100 g/t of feed of XTRACT® (XT hot group) were evaluated by comparison to a negative control (Normal group) and to a positive control being heat stressed (Hot group) but not supplemented with the additive. All animals had *ad libitum* access to fresh water.

Promising standardized plant extracts




The heat stress treatment (from 21°C to 35°C) had a significant negative influence on the treated groups, showing the validity of the used model.


As expected, feed intake reduced when birds were reared under heat stress conditions. The observed

reduction from d. 21 to d. 35 on average daily intake was around 40%. XTRACT® supplementation had no significant effect on daily feed intakes, exactly as would be observed under normal rearing conditions.

Nevertheless, XTRACT® supplemented at 100 g/t had a significant positive effect on body weight gain from d21 to d35 increasing it by 38%. This result, not being a consequence of a higher intake, shows the importance of gut protection, so that the birds can assimilate more efficiently the reduced quantity of ingested nutrients. XTRACT® has the capacity to protect gut epithelium and to reduce its permeability.



The optimal gut protection combined with increased digestive secretions (also observed in normal conditions) directly resulted in limiting performance losses. Heat stressed birds being protected by XTRACT® could achieve more normal FCR level (2.61 vs. 3.29, representing 18 % reduction) during this stressful period.



In this specific trial, the synergy between the 3 ingredients revealed better results than ingredients alone, showing the importance of the mode of actions management.

Because of the natural healthy properties of protectors, some naturally occurring ingredients from plants and spices can protect animals from heat stress and help breeders minimize the economic impact of such extreme conditions. The strategy used in broilers by the Swiss company is an extension of similar positive experiences in ruminants using other plant extract combinations. XTRACT® 6930 can be supplemented to birds through feed but is also available as a soluble instant form in order to be flexibly distributed through drinking water.



Want to learn more ? Contact our expert

Jennifer MAURIN
Business Development Manager for Bioactives
(XTRACT® & NEX®)
Tel: +41 22 929 84 52 | Mob: +41 79 915 6930
jennifer.maurin@pancosma.ch

Pancosma is a global pioneer in developing, manufacturing and distributing a wide range of innovative feed additives. With more than 70 years of experience and real know-how, the company based in Geneva, Switzerland, is present in 75 countries worldwide

www.pancosma.com | Follow us 